

Details of Mittag-Leffler random variate generation

Peter Straka

2017-11-08

First type Mittag-Leffler distribution

Random variate generation

For the efficient generation of random variates, we use the following useful fact (see e.g. Theorem 19.1 in Haubold, Mathai, and Saxena (2011)): A standard α -Mittag-Leffler random variable Y has the representation:

$$Y \stackrel{d}{=} X^{1/\alpha} Z$$

where X is standard exponentially distributed, Z is α -stable with Laplace Transform

$$\mathbf{E}[\exp(-sZ)] = \exp(-s^\alpha),$$

X and Z are independent, and $\stackrel{d}{=}$ means equality in distribution.

Generating X

```
n <- 5
x <- rexp(n)
```

Generating Z

To generate such random variates Z , we use

```
a <- 0.8
sigma <- (cos(pi*a/2))^(1/a)
z <- stabledist::rstable(n = n, alpha = a, beta = 1, gamma = sigma, delta = 0, pm = 1)
```

Below are the details of the calculation. We use the parametrization of the stable distribution by Samorodnitsky and Taqqu (1994) as it has become standard. For $\alpha \in (0, 1)$ and $\alpha \in (1, 2)$,

$$\mathbf{E}[\exp(itZ)] = \exp \left\{ -\sigma^\alpha |t|^\alpha \left[1 - i\beta \operatorname{sgn} t \tan \frac{\pi\alpha}{2} \right] + iat \right\}$$

As in Meerschaert and Scheffler (2001), Equation (7.28), set

$$\sigma^\alpha = CT(1 - \alpha) \cos \frac{\pi\alpha}{2},$$

for some constant $C > 0$, set $\beta = 1$, set $a = 0$, and the log-characteristic function becomes

$$-C \frac{\Gamma(2-\alpha)}{1-\alpha} \cos \frac{\pi\alpha}{2} |t|^\alpha \left[1 - i \operatorname{sgn}(t) \tan \frac{\pi\alpha}{2} \right] \quad (1)$$

$$= -C\Gamma(1-\alpha)|t|^\alpha \left[\cos \frac{\pi\alpha}{2} - i \operatorname{sgn}(t) \sin \frac{\pi\alpha}{2} \right] \quad (2)$$

$$= -C\Gamma(1-\alpha)|t|^\alpha (\exp(-i \operatorname{sgn}(t)\pi/2))^\alpha \quad (3)$$

$$= -C\Gamma(1-\alpha)(-i|t|\operatorname{sgn}(t))^\alpha \quad (4)$$

$$= -C\Gamma(1-\alpha)(-it)^\alpha \quad (5)$$

Setting $t = is$ recovers the Laplace transform, and to match the Laplace transform $\exp(-s^\alpha)$ of Z , it is necessary that $C\Gamma(1-\alpha) = 1$. But then $\sigma^\alpha = \cos(\pi\alpha/2)$, and we see that

$$Y \sim S(\alpha, \beta, \sigma, a) = S(\alpha, 1, \cos(\pi\alpha/2)^{1/\alpha}, 0)$$

Generating Y

```
y <- x^(1/a) * z
y
```

```
## [1] 0.6494067 1.7803382 0.3983979 3.3911503 0.7115620
```

References

- Haubold, H.J., A. M. Mathai, and R. K. Saxena. 2011. "Mittag-Leffler Functions and Their Applications." *J. Appl. Math.* 2011: 1–51. doi:10.1155/2011/298628.
- Meerschaert, Mark M, and Hans-Peter Scheffler. 2001. *Limit Distributions for Sums of Independent Random Vectors: Heavy Tails in Theory and Practice*. Book. First. New York: Wiley-Interscience.
- Samorodnitsky, Gennady, and Murad S Taqqu. 1994. *Stable Non-Gaussian Random Processes: Stochastic Models with Infinite Variance*. Stochastic Modeling. London: Chapman Hall.